



CHROMATOGRAPHY TECHNOLOGY

BLOOD TYPING TEST - CROSSMATCH TEST
DIRECT ANTIGLOBULIN TEST (COOMBS)



BLOOD TYPING TEST

Canine



QuickTest BT DEA 1

Ref. : QT-BT-C

Determination

DEA 1 antigen

Specificity

Anti-DEA 1 monoclonal antibody

Summary

In dogs, the determination of DEA 1 antigen is **strongly recommended before any blood transfusion** to avoid a potent alloantibody response against the DEA 1 antigen.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

1 individual test with all material

PROTOCOL SUMMARY

 **5 MINUTES** procedure

1 DROP of whole blood,
3 DROPS of buffer



SCAN QR CODE :
Movie procedure -
QuickTest BT Canine

Feline



QuickTest BT A+B

Ref. : QT-BT-F

Determination

A/B/AB antigens

Specificity

Anti-A monoclonal antibody
Anti-B monoclonal antibody

Summary

In cats, the presence of naturally occurring alloantibodies in type A and in type B cats requires that **blood typing must be performed prior to blood transfusion** to avoid an acute hemolytic transfusion reaction.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

1 individual test with all material

PROTOCOL SUMMARY

 **5 MINUTES** procedure

1 DROP of whole blood,
3 DROPS of buffer



SCAN QR CODE :
Movie procedure -
QuickTest BT Feline

Equine



QuickTest BT Ca

Ref. : QT-BT-ECa

Determination

Ca antigen

Specificity

Anti-Ca monoclonal antibody

Summary

In horses, there are 7 main blood group systems (A,C,D,K,P,Q,U) with greater than 30 red blood cell factors. Of these, **Ca seems to be one of the most immunogenic** (causing allo-immunization) **antigens**.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

1 individual test with all material

PROTOCOL SUMMARY

 **5 MINUTES** procedure

1 DROP of whole blood,
3 DROPS of buffer

BLOOD TYPING TEST

Canine



Easy QuickTest BT DEA 1

Ref. : EQT-BT-C

Determination

DEA 1 antigen

Specificity

Anti-DEA 1 monoclonal antibody

Summary

In dogs, the determination of DEA 1 antigen is **strongly recommended before any blood transfusion** to avoid a potent alloantibody response against the DEA 1 antigen.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

1 individual test with all material

Feline



Easy QuickTest BT A+B

Ref. : EQT-BT-F

Determination

A / B / AB antigens

Specificity

Anti-A monoclonal antibody
Anti-B monoclonal antibody

Summary

In cats, the presence of naturally occurring alloantibodies in type A and in type B cats requires that **blood typing must be performed prior to blood transfusion** to avoid an acute hemolytic transfusion reaction.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

1 individual test with all material

PROTOCOL SUMMARY

 **5 MINUTES** procedure

1 DROP of whole blood,
3 DROPS of buffer

SCAN QR CODE :
Movie procedure -
Easy QuickTest BT Canine



PROTOCOL SUMMARY

 **5 MINUTES** procedure

1 DROP of whole blood,
3 DROPS of buffer

SCAN QR CODE :
Movie procedure -
Easy QuickTest BT Feline



BLOOD TYPING TEST

Canine



LabTest BT DEA 1

Ref. : LT-BT-C

Determination

DEA 1 antigen

Specificity

Anti-DEA 1 monoclonal antibody

Summary

In dogs, the determination of DEA 1 antigen is **strongly recommended before any blood transfusion** to avoid a potent alloantibody response against this DEA 1 antigen.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

20 tests per box
Additional material required

Feline



LabTest BT A+B

Ref. : LT-BT-F

Determination

A/B/AB antigens

Specificity

Anti-A monoclonal antibody
Anti-B monoclonal antibody

Summary

In cats, the presence of naturally occurring alloantibodies in type A and in type B cats requires that **blood typing must be performed prior to blood transfusion** to avoid an acute hemolytic transfusion reaction.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life and storage

18 months at room temperature

Presentation

20 tests per box
Additional material required

PROTOCOL SUMMARY



5 MINUTES procedure

1 DROP of whole blood,
3 DROPS of buffer

SCAN QR CODE :
Movie procedure -
LabTest BT Canine



PROTOCOL SUMMARY



5 MINUTES procedure

1 DROP of whole blood,
3 DROPS of buffer

SCAN QR CODE :
Movie procedure -
LabTest BT Feline



CROSSMATCH TEST (XM)

Canine



QuickTest XM

Ref. : QT-XM-C

Detection

Canine allo-antibodies

Specificity

Canine IgG/IgM/C3 Anti-globulin

Summary

Our Canine Crossmatch Test will allow you to pick up incompatibilities across **ALL** canine blood groups : DEA 1, 3, 4, 5, 7, DAL... For safe transfusion, it is mandatory to perform both blood typing and crossmatch testing.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life & storage

18 months at room temperature

Presentation

1 individual test with all material

Other packaging

Lab Test XM

5 tests per tube with all material

Feline



QuickTest XM

Ref. : QT-XM-F

Detection

Feline allo-antibodies

Specificity

Feline IgG/IgM/C3 Anti-globulin

Summary

Our Feline Crossmatch Test will allow you to pick-up incompatibilities across **all** feline blood groups : A, B, AB, Mik and other unknown antigens. For safe transfusion, it is mandatory to perform both blood typing and crossmatch testing.

Sample type

Anticoagulated whole blood (EDTA/ACD/CPD)
Do not use Heparin

Shelf life & storage

18 months at room temperature

Presentation

1 individual test with all material

Other packaging

Lab Test XM

5 tests per tube with all material

PROTOCOL SUMMARY

 **25 MINUTES** procedure

1 DROP of pRBCs
3 DROPS of plasma
10 MINUTES incubation
WASHING STEP PROCEDURE NEEDED

SCAN
QR CODE :
Procedure -
QuickTest
XM Canine



PROTOCOL SUMMARY

 **25 MINUTES** procedure

1 DROP of pRBCs
3 DROPS of plasma
10 MINUTES incubation
WASHING STEP PROCEDURE NEEDED

SCAN
QR CODE :
Procedure -
QuickTest
XM Feline



Discover our new app

Free App available
on all mobile phones

Archivable results with
PDF report by email

Instant result using AI and
operator confirmation




Scientific assistance
within 24 hours

In-App video
instructions of
test procedure

Troubleshooting
documentations

Please, scan the
QR code to open
the application on
your smartphone



ADDRESS

11-13 Rue des Aulnes,
69760 Limonest
France

PHONE

+33 478 380 239

E-MAIL

contact@alvedia.com

WEBSITE

www.alvedia.com

